


SOLUTIONS

1.

$$F(x) = x\sqrt{6-x}$$


 (a) What is the domain of $F(x)$? $(-\infty, 6]$

(b) Find the intervals of increase or decrease and critical numbers.

(c) Find the intervals of concavity and the inflection points.

(d) Sketch the graph.

increase: $(-\infty, 4]$
 decrease: $[4, 6]$
 crit. numbers: $x=4, 6$
 concave down $(-\infty, 6)$
 never concave up
 no inflection points

$$F'(x) = 1 \cdot \sqrt{6-x} + x \cdot \frac{1}{2}(6-x)^{-\frac{1}{2}}(-1)$$

$$= (6-x)^{-\frac{1}{2}} ((6-x) - \frac{x}{2}) \quad \leftarrow \text{factor } \underline{\text{lowest power}}$$

$$= \frac{6 - \frac{3}{2}x}{\sqrt{6-x}}$$

$$F''(x) = \frac{(-\frac{3}{2})(6-x)^{\frac{1}{2}} - (6 - \frac{3}{2}x)^{\frac{1}{2}}(6-x)^{-\frac{1}{2}}(-1)}{6-x}$$

$$= \frac{(6-x)^{-\frac{1}{2}} [-\frac{3}{2}(6-x) + \frac{1}{2}(6 - \frac{3}{2}x)]}{6-x} = \frac{\frac{3}{4}x - 6}{(6-x)^{\frac{3}{2}}}$$

x	F	F'	F''
0	0	0	
4	$4\sqrt{2}$	0	
6	0	d.n.e.	

2.

$$f(t) = t^{4/5}(t-4)^2$$

 (a) What is the domain of $f(t)$? $(-\infty, \infty)$

 (b) Find $f'(t)$. What is its domain? formula below; $(-\infty, 0) \cup (0, \infty)$

(c) Find all the critical numbers.

$$C = 0, \frac{8}{7}, 4$$


$$f'(t) = \frac{4}{5}t^{-\frac{1}{5}}(t-4)^2 + t^{\frac{4}{5}}2(t-4)$$

factor lowest powers

$$= t^{-\frac{1}{5}}(t-4) \left[\frac{4}{5}(t-4) + 2t \right]$$

$$= \frac{(t-4)(\frac{14}{5}t - \frac{16}{5})}{t^{\frac{1}{5}}}$$

$\therefore f'(c)=0$ at $c=4, \frac{8}{7}$
 $f'(c)$ d.n.e. at $c=0$



3.

$$g(x) = \frac{e^x}{1 - e^x}$$

- (a) What is the domain of $g(x)$? $X \neq 0$ or $(-\infty, 0) \cup (0, \infty)$ $\therefore x=0$ is vertical $y=0$ is hor.
- (b) Find the horizontal and vertical asymptotes. $\lim_{x \rightarrow 0^-} g(x) = +\infty$, $\lim_{x \rightarrow 0^+} g(x) = -1$, $\lim_{x \rightarrow \infty} g(x) = 1$ \leftarrow §4.4 helps with kind of limit you can do it anyway! $y=1$ is hor. $y=-1$ is hor.
- (c) Find the intervals of increase or decrease and critical numbers. { No critical #s
Increase: $(-\infty, 0) \cup (0, \infty)$
Never decreasing }
- (d) Find the intervals of concavity and the inflection points. Concave up on $(-\infty, 0)$, concave down on $(0, \infty)$, no infl. pts
- (e) Sketch the graph.

$$g'(x) = \frac{e^x(1-e^x) - e^x(-e^x)}{(1-e^x)^2} = \frac{e^x}{(1-e^x)^2} \therefore g'(c) = 0$$

$$g''(x) = \frac{e^x(1-e^x)^2 - e^x 2(1-e^x)(-e^x)}{(1-e^x)^4} = \frac{e^x(1-e^x) + 2e^x e^x}{(1-e^x)^3}$$

$$= \frac{e^x(1+e^x)}{(1-e^x)^3} \therefore g''(c) = 0 \text{ has no solutions}$$

and $g'(x) > 0$ if $x < 0$

$g'(x) < 0$ if $x > 0$

x	g	g'	g''
-1	+0.6	+	+
0	d.n.e.	d.n.e.	d.n.e.
1	-1.6	+	-

